

[CLICK HERE TO JUMP TO USE DIRECTIONS](#)



MECOMEC® 2.5 TURF HERBICIDE

For Control of Clover and other Broadleaf Weeds in Golf Greens, Fairways, and Lawns.

ACTIVE INGREDIENT:

Potassium salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid 15.55%

INERT INGREDIENTS: 84.45%
TOTAL 100.00%

THIS PRODUCT CONTAINS:

Equivalent to 13.20% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent or contains 1.16 lbs. MCP-p per gallon.

Contains single isomer form of Mecoprop-p.

KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional Precautionary Statements and First Aid.

KEEP FROM FREEZING



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Avoid contact with skin, eyes and clothing. Wear long pants, long-sleeved shirt, shoes and gloves when mixing, loading or applying this product. Wear face shield or goggles when mixing, loading or applying this product. Wash hands after use.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. <p style="text-align: right;"><i>(continued on next column)</i></p>

FIRST AID (cont.)

If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
--------------------	---

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as MCP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not reenter or permit workers to reenter treated area until sprays have dried or dusts have settled.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing. Do not store open containers near seeds, feeds, fertilizer or pesticides including insecticides, fungicides and herbicides other than those containing 2,4-D or brush killers.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

MECOMEC® 2.5 Turf Herbicide will not injure bentgrass, established lawns or turfgrass when used as directed. Controls mouseear chickweed, clover, plantains, black medic and knotweed in turfgrass.

Apply when weeds are young and actively growing. Do not apply to bentgrasses in the heat of the summer. Do not mow for several days before or after treatment. Do not water for one day following treatment.

Do not allow spray mist, visible or invisible vapors, or air containing mist or vapors to contact plants, other than those being sprayed. If 2,4-D or MCP-p susceptible plants are growing within a distance of 1/2 mile from the area being sprayed, make application only when there is a light breeze away from the susceptible plants. Spray drift can be reduced by using high volume sprays under low pressure, coarse droplet nozzles and boom nozzle drops. Susceptible crops include grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, cauliflower and cabbage, boxelder trees, as well as many other field, vegetables, flowers, ornamental and fruit crops. Do not contaminate streams, lakes, ponds, irrigation water, or water used for livestock or domestic purposes.

Sprayers which have contained MCP-p should not be used to apply other pesticides to 2,4-D or MCP-p susceptible crops.

Do not apply this product through any type of irrigation system.

ESTABLISHED GRASSES (Fairways and Lawns):

For broadcast applications and spot treatments: Apply 1 1/2-2 1/4 fluid ounces of product per 1,000 square feet of turfgrass with spray volumes of 0.5-4.0 gallons per 1,000 square feet. Or, the equivalent dosage rate is 4 to 6 pints of product with spray volumes ranging from 22-174 gallons per acre.

NOTE: If dandelions are a problem, add 2,4-D at the recommended rate except on bentgrasses where the rate of 2,4-D should not exceed 4 oz. acid equivalent per acre.

ESTABLISHED GREENS:

For broadcast applications and spot treatments to control mouseear chickweed, black medic, knotweed, and plantains: Apply 1½ fluid ounces of product per 1,000 square feet of turfgrass with spray volumes ranging from 0.5-4.0 gallons per 1,000 square feet. Or, the equivalent dosage rate is 4 pints of product with spray volumes ranging from 22-174 gallons per acre.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

MECOMEC® is a registered trademark of PBI/Gordon Corporation.

892/6-2004 AP062703
EPA REG. NO. 33955-483-2217

Distributed By



An Employee-Owned Company

**1217 West 12th Street
Kansas City, Missouri 64101**

www.pbigordon.com

SP
L
A
B
E
L
S
P
E
C
I
M
E
N